

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/318744800>

Biodiversity conservation in urban environments: a review on the importance of spatial patterning of landscapes

Conference Paper · July 2017

CITATION

1

READS

1,261

3 authors:



[Amin Rastandeh](#)

University of South Dakota

15 PUBLICATIONS 31 CITATIONS

SEE PROFILE



[Daniel Brown](#)

Victoria University of Wellington

9 PUBLICATIONS 15 CITATIONS

SEE PROFILE



[Maibritt Pedersen Zari](#)

Victoria University of Wellington

59 PUBLICATIONS 416 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Regenerative Cities - Ecosystems Services Analysis [View project](#)



Teaching and education [View project](#)

Please cite this paper as follows:

Rastandeh, A., Brown, D., K., Pedersen Zari, M. (2017). Biodiversity conservation in urban environments: a review on the importance of spatial patterning of landscapes. *Ecocity World Summit*, 12-14 July. Melbourne, Australia.

Biodiversity conservation in urban environments: a review on the importance of spatial patterning of landscapes

Amin Rastandeh, Daniel K Brown, Maibritt Pedersen Zari
Victoria University of Wellington, Wellington, New Zealand

Abstract

It has been well established that biodiversity plays an irreplaceable role in ensuring the quality of human life through supporting ecosystem functions and services. As more and more people prefer to live in cities worldwide, biodiversity loss in urban environments is being increasingly reported more than ever before. This, in turn, may have a negative influence on the quality of human life in an urbanising world. Global research shows that the abundance and richness of fauna in urban environments depends, to a large extent, on the spatial patterning of different patches of urban vegetation such as urban forests, woodlands, parks, and gardens.

The principal aim of the research is to provide a coherent picture of the importance of spatial patterning and spatial ecology of wildlife species in urban environments. Based upon empirical data from North America, Latin America, Europe, Asia, Africa, and Oceania, the research involves a systematic review of international peer-reviewed publications relating to the connection between biodiversity and the composition and configuration of urban wildlife habitats. This review reveals the most important components of landscape pattern that contribute to the abundance and richness of urban wildlife species.

Ultimately, the results provide a deeper understanding of the strategic importance of spatial dimensions of landscape planning and management, in support of biodiversity conservation in landscapes that have already been widely affected by anthropogenic development. Importantly, the findings provide a set of spatially-explicit recommendations that can be strategically applied in urban landscape architecture and land use planning disciplines to help ensure that urban biodiversity is maintained in an era of climate change and rapid urbanisation.

Keywords:

Urban biodiversity; wildlife habitats; spatially-explicit landscape patterns; spatial ecology; landscape architecture

1. Introduction

The importance of biodiversity can be linked to the role of multiple services provided by the functioning of ecosystems. There is a positive relationship between biodiversity and ecosystem functions and services. Balvanera et al. (2006) analysed more than 400 measures of biodiversity effects on ecosystem services and suggested that biodiversity has positive effects on services provided by ecosystems under study. The meta-analysis by Cardinale et al. (2006) shows that biodiversity loss has negative effects on ecosystem functions in many different ways, and the connection between biodiversity and the functioning of ecosystems has been affirmed by others (Hector and Bagchi, 2007; Duffy et al., 2007; Duffy, 2008; Isbell et al., 2011; Hooper et al., 2012; Pasari et al., 2013; Tilman et al., 2014; Lefcheck et al., 2015). Ecosystem services, in turn, support different aspects of the quality of human life (Tzoulas et al., 2007; Pedersen Zari, 2015). These services have been classified as provisioning, regulating, cultural, and supporting services (Millennium Ecosystem Assessment, 2005). The first three services have direct effects on the quality of human life while the fourth supports other services. In an era of rapid urbanisation (Bashford, 2014) and climate change (IPCC, 1995), urban biodiversity is critically important to support the citizens' quality of life. Despite this, biodiversity is threatened by both urbanisation and climate change worldwide (McKinney, 2002; Parmesan, 2006). Grimm et al. (2008), for example, blame biodiversity loss on rapid urbanisation and climate change. They suggest, however, that both problems and solutions can be found in cities. Approximately 52% of the world's population live in cities while in some regions of the world this figure is higher than 80% (United Nations, 2008). This trend is estimated to continue over the coming decades (United Nations, 2014); more people are

expected to live in urban environments and, consequently, more houses and infrastructures are inevitably required in order to respond to the increasing urban population. If managed inappropriately, urban development coupled with climate change impacts is likely to cause more widespread biodiversity loss and this, in turn, may affect the healthy functioning of ecosystems and accordingly the quality of human life in cities.

In response to this challenge, some researchers suggest that composition and configuration of patches of vegetation should be considered as a key vehicle in order for the landscape architecture discipline to respond to urban biodiversity loss. Spatial patterning of patches of urban vegetation is thought to hold the key to providing relatively suitable conditions for urban wildlife to remain in cities and support ecosystem services despite ongoing pressures on wildlife habitats imposed by urban development and climate change. The connection between landscape pattern and urban biodiversity can be interpreted under the pattern-process-relationship model in landscape ecology (Forman and Godron, 1986; Turner, 1989; Forman, 1995; Bell, 2001). According to this concept, patterns affect processes and vice versa at different scales (Forman, 1995; Botequilha Leitaó and Ahern, 2002). Thus, change in each component of landscape pattern can affect urban biodiversity either directly or indirectly. Therefore, it seems imperative for landscape architecture researchers and practitioners to develop knowledge of what the most important components of landscape pattern composition and configuration are and how spatial patterning of urban green spaces may affect the presence, abundance and richness of urban wildlife species.

The principal aim of this research is to provide a coherent picture of the importance of spatial patterning and spatial ecology of wildlife species in urban environments in order for landscape architecture researchers and practitioners to build-up a deeper understanding of the most important components of landscape pattern that contribute to biodiversity in cities. This will help cities to support biodiversity where landscapes and wildlife habitats have already been widely affected.

2. Methodology

From November 2015 to February 2017, international literature was comprehensively reviewed using a wide range of available databases including ISI Web of Science, Science Direct, Springer Links, Scopus, Wiley Online Library, and Pro Quest Central. The aim was to address the following issues:

- (1) To build up an in-depth understanding of pattern process relationship in landscape ecology science using seminal publications in the field;
- (2) To develop an understanding of the role of landscape pattern composition and configuration in affecting wildlife species richness and abundance at the urban scale;
- (3) To identify the most influential components of landscape pattern constructing and characterising urban wildlife habitats and their influence on the presence, richness and abundance of wildlife species.

A wide range of relevant keywords was used to retrieve a large number of publications. Keywords included, but were not limited to, *urban biodiversity*, *urban wildlife*, *urban bird diversity*, *urban wildlife management*, *urban zoology*, *urban avian diversity*, and *urban avifauna*. Keywords were searched both separately and accumulatively to find the most relevant published data. Publications retrieved were filtered to gather a limited number of the most preeminent and reliable ones. Google Scholar Citation Index and Journals' Impact Factors were utilised to examine the quality of publications. Both positive and negative citations were checked for the latest group of publications to ensure that the collection of the selected publications has the highest degree of validity and reliability. Next, the literature review was fulfilled in two stages as follows:

Stage 1: Given that landscape ecology science confirms that *pattern process* relationship is valid in various landscapes and different scales (Turner, 1989; Forman, 1995; Bell, 2001), seminal publications of international reputation on the connection between biodiversity and landscape pattern composition and configuration were explored irrespective of scale of the studies (i.e. urban or regional).

Stage 2: Highly preeminent empirical research on the connection between biodiversity and landscape pattern composition and configuration at the urban scale was reviewed. Geographical diversity of the reviewed research was constantly controlled to ensure that the review considers the global scale and includes the diversity of climatic zones. Four criteria were taken into particular consideration in order to find the most relevant published data in this stage:

- (1) Peer-reviewed, published from 2000 to the end of February 2017, written in English and indexed on online databases;

- (2) Empirical-based, focused on the connection between landscape pattern composition and configuration and urban wildlife species richness and abundance on an urban scale;
- (3) Received international attention, including a reasonable number of positive citations recorded on Google Scholar;
- (4) Represented a local and/or regional picture of the current issues in the region under study.

Ultimately, forty-one urban scale studies comprising empirical research in Africa (n = 2), Asia (n = 12), Europe (n = 9), Latin America (n = 6), North America (n = 5), and Oceania (n = 7) were selected for the final review representing a global perspective of the topic under study (Figure 1).




Figure 1. The multi-stage/criteria process undertaken to find the most reliable and relevant publications relating to the relationship between urban biodiversity and landscape pattern composition and configuration.

3. Results

3.1. Components of landscape pattern

Conceptual models (e.g. Island Biogeography Model (Mac Arthur and Wilson, 1967), Source-Sink Model (Pulliam, 1988), Patch-Corridor-Matrix Model, Forman, 1995), and the like) provide evidence that biodiversity is profoundly dependent on landscape pattern composition and configuration (Forman and Godron 1981; Forman and Godron 1986; Bridgewater, 1988; Turner, 1989; Soulé, 1991; Noss and Cooperrider, 1994; Forman, 1995; Murcia 1995; Dramstad et al., 1996; Collinge, 1996; Forman and Collinge, 1997; Savard, 2000; Noss, 2001; McGarigal and Cushman, 2002; Environmental Law Institute, 2003; Alberti, 2005; Botequilha Leitao et al., 2006; Farina, 2006; Opdam et al., 2006; Fischer et al., 2009; Haddad, 2009; Moorcroft, 2009; Heller and Zavaleta, 2009; Morrison et al., 2012; Walz and Syrbe, 2013). Landscape pattern is constructed and characterised by a number of components. From the literature reviewed, the most important components of landscape pattern affecting biodiversity are as follows:

- (1) Indigenousness;
- (2) Land cover heterogeneity;
- (3) Land surface perviousness;

- (4) Patch size;
- (5) Connectivity and proximity;
- (6) Edge density and contrast;
- (7) Landform diversity;
- (8) Shape complexity.

Taken together, the above-mentioned components of landscape pattern construct spatial characteristics of wildlife habitats, both in urban or non-urban environments. Change in any component of landscape pattern will have different implications for wildlife species richness and abundance. The components and their relationships with biodiversity are discussed in the following sections.

3.1.1. *Indigenesness*

Indigenesness is defined as the state of being indigenous (i.e. native) in terms of land cover type classes present in a landscape. Indigenous plants have an irreplaceable role in biodiversity – and thereby long-term ecosystem integrity in cities (Aronson et al., 2014; Rastandeh, et al., 2017). Forman (1995) suggests maintaining patches of indigenous vegetation, as small *bits of nature*, within human modified landscapes because of the services they provide as habitat and/or stepping stones for a wide range of wildlife species. Fischer et al. (2009) state that the overall level of biodiversity depends on the total number of indigenous plants present in a given landscape. The positive relationship between the percentages of indigenous vegetation and the extinction rate of indigenous plants has been confirmed in cities (Hahs et al., 2009).

3.1.2. *Land cover heterogeneity*

Land cover heterogeneity is defined as diversity of different land cover type classes in patch or landscape levels. The number of habitat types is positively related to biodiversity across landscape (Forman and Godron, 1986). Dramstad et al. (1996) argue that land cover diversity can contribute to biodiversity in different scales. Nielsen et al. (2013) reviewed 62 empirical research studies from 25 countries and pointed out that land cover heterogeneity may be the most important factor supporting urban biodiversity. In large scale, Fischer et al. (2009) affirmed that since species differ in their habitat requirements, diversity in land cover types in the same landscape can underpin suitable conditions for the presence and survival of different types of species. Botequilha Leitaó et al. (2006) argue that some key ecological functions are affected by land cover diversity. They explain how richer diversity in forest and/or grassland land cover types can cause greater biodiversity. Reside et al. (2014) suggest that a wide range of habitats are required to maintain the long-term evolutionary process of species present across the landscape. Land cover heterogeneity can also give rise to a larger number of eco-tones where the number of wildlife species is thought to be high (Duelli, 1997).

3.1.3. *Land surface perviousness*

Land surface perviousness is defined as the ability of a particular land cover type or landscape to absorb run-off caused by rainfall, or sequester and store carbon dioxide in soil or vegetation. Some studies show that there is a positive relationship between land surface perviousness and bird diversity and abundance in urban landscapes. For example, Fernández-Juricic and Jokimaki (2001) demonstrate the negative correlation between bird species richness and paved ground within urban parks in Madrid, Spain. A study of three cities in Switzerland by Fontana et al. (2011) revealed that the percentage of land dominated by trees is the most important variable enhancing bird diversity in urban landscapes. A study of bellbird occupancy in Christchurch, New Zealand showed that the chance of bellbird presence is higher in unpaved urban surfaces covered by indigenous plants (MacLeod et al., 2012).

3.1.4. *Patch size*

Patch size is defined as the total area of a particular land cover type on a patch or landscape level. Other variables being equal, an individual large habitat can support more species because it is regarded as a large pool of species genes (Forman, 1995; Dramstad et al., 1996; ELI, 2003; Fischer et al., 2009). Large patches of vegetation are likely to contain more diverse species, thereby providing more widespread ecosystem services. Forman and Godron (1981) suggest that the size of patches present in a landscape affects

productivity, nutrient and water flux, and species dynamics and, therefore, can be considered as an important indicator of biodiversity. Based upon MacArthur and Wilson (1967), they argued that when habitat diversity (i.e. heterogeneity) is equal in two patches of vegetation, the larger patch typically contains more species (cf. Forman et al., 1976; Forman and Godron, 1981).

3.1.5. Connectivity and proximity

Connectivity is defined as the spatial distance between patches of a particular land cover type. Spatial connectivity is believed to be vital to biodiversity (Naveh, 1994; Forman, 1995; Bennett, 1999). Spatial connectivity between patches of vegetation can facilitate the process of colonisation and, thereby, reduce the chance of plant species extinction (Shaffer, 1981; Honnay et al., 2003). The concept of planning urban and regional greenways that emerged in landscape architecture and regional planning (Ahern, 1995; Fabos, 2004; Toccolini, 2006; Turner, 2006; Walmsley, 2006; Mason, 2007; Teng et al., 2011; Palmisano et al., 2016) has been, to a large extent, a response to this ecological necessity. Some ecological processes and species, but not all, benefit from connectivity between patches including movement of species between habitats and the flux of energy and nutrients (Fischer et al., 2009). Connected patches in urban environments provide buffered corridors for species to migrate from one habitat to another to find food and shelter without facing unpleasant climatic conditions and urbanisation effects. Conversely, some researchers argue that connectivity can concurrently facilitate the dispersal of weeds and pests (Barnes, 2000; Botequilha Leitao et al., 2006; Sullivan et al., 2009).

3.1.6. Edge density and contrast

Edge density is defined as the perimeter of a patch of particular land cover type exposed to other land cover types. Edge contrast is also defined as dissimilarity between adjacent land cover types. Landscape fragmentation increases edge density (Andren, 1994). Both edge density (i.e. length) and edge contrast can affect urban biodiversity. As edge density increases, the patch interaction with its surroundings increases either positively or negatively (Dramstad et al., 1996). In a New Zealand context, Young and Mitchell (1984) and Davies-Colley et al. (2000) revealed that climatic edge effects are considerably reduced within c. 50 m and c. 40 m from the patch boundary, respectively. The former threshold has been suggested by Meurk and Swaffield (2000) and Meurk and Hall (2006) to be regarded as a basis for edge effect analysis in New Zealand urban landscapes. This threshold has also been suggested by Murica (1995) after reviewing a comprehensive number of relevant publications, as well. In Australia, a 50 m edge width was recommended to design buffer zones for highly vulnerable large-size body species such as koala (Port Stephens Council, 2002). As wind and sun exposure widen edge effects (Dramstad et al., 1996), it is not, however, correct to assume that the width of edge effect remains the same around a patch.

3.1.7. Landform diversity

Landform diversity is defined as the diversity of elevations, slopes, and aspects. Landform diversity has been strongly suggested to safeguard biodiversity in the face of extreme events (Markham et al., 1993; Pernetta et al., 1994; Halpin, 1997). Research shows that temperature differences between south- and north-facing slopes can be considerable, ranging from 8°C to 12°C (Rorison et al., 1986; Ackerly et al., 2010). Landform diversity provides climate heterogeneity, and thereby safeguards more diverse species over time due to the diversity in environmental variables such as temperature, moisture and soil type in a heterogeneous topography (Noss, 2001; Fischer et al., 2009; Dobrowski et al., 2011; Reside et al., 2014). On an urban scale, diverse topography potentially provides diverse habitats for a wide range of urban species. On a fine scale (i.e. small urban parklands), however, research shows that landform diversity is less important to urban wildlife species (Cornelis and Hermy, 2004).

3.1.8. Shape complexity

Shape complexity is defined as the degree to which a wildlife habitat is dissimilar to a circle-shaped pattern. The proportion between edge-core areas in a given patch depends on patch shape affecting biodiversity (Noss and Cooperrider, 1994; Dramstad et al., 1996); however, the nature of impact may differ depending on the species under study. O'Neill et al. (1988) and Honnay et al. (1999) showed that patches with irregular shape contribute to richer plant diversity. At the same time, however, a wide range of animals may prefer

compact shapes to protect themselves from the edge effects (Forman, 1995) because compact shapes decrease the odds of the penetration of negative environmental effects into the patch. Botequilha Leitaó et al. (2006) depicted the relationships between patch shape, core area, and biodiversity and discussed how some characteristics are expected to occur in landscape when patch shape complexity increases and patch core area consequently decreases. They enumerated a range of events including an increase in the rate of evapotranspiration along patch edges exposed to sunlight and wind, an increase in the numbers and populations of exotic and predator species (and at the same time, a decrease in the presence of rare plant species that require specific patch interior conditions), a sharp reduction in the populations of ground-nesting songbirds due to predation pressure, and a decrease in storm-water storage in remnant patches of vegetation (Botequilha Leitaó et al., 2006).

3.2. Empirical evidence in urban environments

Recently more attention has been paid to biodiversity and spatial ecology of wildlife species in urban environments (Müller and Kamada, 2011). The results of the literature review show an increasing trend in the number of publications on the relationship between spatial ecology of wildlife species and spatial composition and configuration of patches of vegetation in urban environments from 2001 to February 2017. Empirical evidence worldwide within a spectrum of different climatic zones (Peel et al., 2007) verifies that the presence, richness and abundance of urban wildlife species rely profoundly on spatial pattern of wildlife habitats scattered across urban environments. The selected publications are testimony to the role of the eight most important components of landscape pattern (indigenesness, land cover heterogeneity, land surface perviousness, patch size, connectivity and proximity, edge density and contrast, landform diversity, and shape complexity) in affecting a wide range of urban wildlife species including various butterflies, birds, reptiles, and mammals. The results derived from this stage of the literature review have been summarised to depict a global picture of the preeminent publications in peer-reviewed journals of international repute (Table 1).

Table 1: Selected international peer-reviewed publications based upon empirical data and field surveys from 2000 to February 2017 addressing the role of landscape pattern in affecting wildlife species in urban environments. Components involve: (1) indigenouness, (2) land cover heterogeneity, (3) land surface perviousness, (4) patch size, (5) connectivity and proximity, (6) edge density and contrast, (7) landform diversity, and (8) shape complexity. For additional information about climatic zones, see Appendix I.

Continent	City – Urban region	Climatic zone	Species under study	Components affecting species	Reference
North America					
Canada	Southern Ontario	Dfa, Dfb, Dfc	Birds	1, 2, 4	Smith, 2007
	St. Louis	Dfa - Cfa	Birds	2, 4	Oliver et al., 2011
	Boise	BSk	Birds	3	McClure et al., 2015
	Boston	Cfa	Birds	3, 4	Strohbach et al., 2013
	Baltimore	Cfa	Birds	3, 6	Rega et al., 2015
Europe					
Belgium	Flanders	Cfb	Butterflies, birds, amphibians	1, 2, 4	Cornelis and Hermly, 2004
	Prague	Cfb	Birds	1, 2, 5	Ferenc et al., 2014
	Marseille	Csa	Butterflies	5, 8	Lizee et al., 2012
	Halle	Cfb	Butterflies, birds, invertebrates	4, 6, 8	Bräuniger et al., 2010
	Madrid	Csa	Birds	2, 4, 6	Fernández-juricic, 2001
	Valencia	BSk	Birds	2, 3, 4	Murgui, 2009
	Orebro	Dfb	Birds	1, 2, 5	Sandstrom et al., 2006
	Zurich, Lucerne, Lugano	Cfb	Birds	2, 3	Fontana et al., 2011
	100 urban areas	Cfb	Birds	2, 4, 6	Evans et al., 2009
Latin America					
Argentina	Mar del Plata	Cfb	Birds	3, 5, 6	Leveau and Leveau, 2016
	La Paz	Cwb	Birds	1, 3, 7	Villegas and Garitano-Zavala, 2010
	Taubate	Cwa	Birds	1, 2	de Toledo et al., 2012
	Uberlandia	Aw	Invertebrates	1, 5	Pacheco and Vasconcelos, 2007
	Pachuca	Cwb	Birds	2, 4, 6	Carbó-Ramírez and Zuria, 2011
Venezuela	Margarita Island	Aw	Birds	1, 3, 5	Sanz and Caula, 2015
Africa					
South Africa	Cape Town	Csb	Birds	5, 6	Calder et al., 2015
	Cape Town	Csb	Birds	2, 5	Suri et al., 2017

Table 1: continued

Continent	City	Climatic zone	Species under study	Components affecting species	Reference
Asia	China	Dwa	Birds	2, 5	Yuan and Lu, 2016
	Japan	Cfa	Birds	3, 6	Imai and Nakashizuka, 2010
	Hong Kong	Cwa	Birds	4	Zhou and Chu, 2012
	India		Birds	1, 2, 4	Khera et al., 2009
	Israel	Csa	Birds	2, 3, 5	Shwartz et al., 2008
		Csa	Birds	1, 2	Paker et al., 2014
	Malaysia	Af	Butterflies	2, 4, 5	Sing et al., 2016
		Af	Birds	4, 6	Aida et al., 2016
		Af	Birds	1, 3, 4	Jasmani et al., 2017
	Philippines	Aw	Birds	2, 4	Vallejo et al., 2009
	Singapore	Af	Butterflies, birds	1, 3, 6	Chong et al., 2014
	South Korea	Dwa	Birds	2, 4, 5	Kang et al., 2015
Oceania	Australia	Cfa	Reptiles, small mammals	1, 2, 3, 4, 8	Garden et al., 2010
		Cfb	Birds	2, 3, 4	Ikin et al., 2013
		Cfb	Birds, bats	1, 2	Threlfall et al., 2016
		Cfb	Birds	4, 5	Platt and Lill, 2006
	New Zealand	Cfb	Birds	1, 2, 4, 5	vanHeezik et al., 2008
		Cfb	Birds	1, 3, 4	vanHeezik et al., 2013
		Cfb	Birds	1, 3, 5	MacLeod et al., 2012

4. Discussion

4.1. A more realistic approach

Although *making cities as green as possible* seems to contribute to urban biodiversity to some extent, implementation of this strategy may be unrealistic in practice because urban green space development requires enough space (i.e. suitable land) and sufficient funds. Beyond these factors, water shortage resulting from worldwide population growth and global warming is very likely to limit the goal of *making cities as green as possible*, specifically in countries where climate change affects water resources (IPCC, 1995; IPCC, 2014). Therefore, planning for biodiversity conservation in cities may differ in some cases from what planners undertake in natural areas. Knowledge of the spatial ecology of wildlife species, however, may help urban policy makers to make appropriate decisions on not only allocation of land to green spaces, but also spatial design of patches of urban vegetation (i.e. parklands, woodlands, and urban forests) in a way that supports wildlife species in urban environments where land availability is widely limited by socio-economic drivers.

4.2. Areas of conflict

This research also reveals that in terms of spatial patterning of patches of vegetation in urban environments, there are some areas of conflict between what wildlife species require and what humans expect from ecosystem services. First, although indigenous plants are necessary to ensure the continuation of biodiversity in urban environments, some exotic species may be socio-economically more acceptable to be used by land owners for carbon sequestration and storage (Ausseil et al., 2013; Ministry for Primary Industries, 2015; Setälä et al., 2016; McHale et al., 2017). In an era of climate change, conventional strategies may encourage people to plant exotic species to manage greenhouse gasses in cities through carbon sequestration. At the same time, research shows that some exotic flora may provide indigenous fauna with essential food during winters or initiate ecological succession towards indigenous plant communities (Rastandeh et al., 2017). Therefore, use of only indigenous plants in urban environments should not be considered as a clear-cut response to biodiversity loss because cities already contain non-indigenous plants, and these can be beneficial to urban ecosystems.

Second, while a higher rate of shape complexity and accordingly edge density may decrease the quality of wildlife habitats and consequently affect urban biodiversity, it can also reduce urban heat island effects through increasing the cooling effects of urban green spaces. A study of 21 urban parks in Addis Ababa, Ethiopia, for example, revealed that an increase in shape complexity and edge density may increase the park cooling distance – the range within which the cooling effect could be observed (Feyisa et al., 2014). This example is compatible with the results of a study conducted in the city of Aksu, China (Maimaitiyiming et al., 2014) where researchers showed that greater edge density resultant from higher shape complexity can effectively reduce land surface temperatures in urban environments without the need for increasing the total green space area. Similarly, a spatial analysis of land cover composition and configuration in the hot-arid city of Isfahan, Iran concludes that shape complexity (i.e. irregular, elongated, and convoluted configuration with enough core area) is negatively correlated with land surface temperature in urban environments (Asgarian et al., 2015; cf. Zhou et al., 2011; Kong et al., 2014; Park and Cho, 2016). Therefore, while compact patches of vegetation may benefit wildlife species through providing more core area and less edge, elongated ones may be more beneficial for people suffering from urban heat island effects.

Due to the coexistence of humans and wildlife species in cities, some ecosystem services may become contradictory in urban environments. Thus, the question of *high vs. low shape complexity* and consequently *high vs. low edge density* may remain an important issue of concern in landscape architecture research, particularly in the contemporary urban age when urban policy makers seek solutions to adapt cities and people to the local impacts of climate change. Therefore, a middle ground should be sought in order for landscape architecture researchers and practitioners to meet both wildlife species and human requirements. Such paradoxical functioning of ecosystems in non-urban environments is, however, unlikely because in natural landscapes planning is solely focused on wildlife conservation while in urban environments both wildlife species and humans must be taken

into consideration in the process of planning patches of vegetation, and this in itself makes this process more complex and multi-dimensional (Figure 2).




Figure 2. An example of areas of conflict between what wildlife species require and what people expect from spatial patterning of patches of vegetation in urban environments. Although green space area in both cases is equal (A), shape complexity, and thereby edge density, profoundly differs. The compact pattern is suitable for wildlife species to be protected from edge effects (B), whereas the dispersed pattern is more suitable for humans to mitigate urban heat island effects (C). In urban environments, there seems to be a need to seek a middle ground between what humans and wildlife species require in terms of spatial patterning of patches of vegetation.

4.3. A spatially-explicit perspective

Although planning for urban biodiversity is extremely site- and species-specific (Turner, 1989; Botequilha Leitao and Ahern, 2002; Botequilha Leitao et al., 2006), the current literature provides a coherent picture of what wildlife species may require in urban environments to survive. According to

the research results, an ideal spatial pattern for wildlife habitat that ensures long-term presence, richness and abundance of wildlife species – and at the same time considers what humans require in the face of climate change-related issues such as urban heat island effects – can be summarised as follows:

- (1) A combination of patches of different sizes from small to large, mostly connected through green and blue corridors, where possible, as well as a limited number of isolated patches of vegetation as urban wildlife refugia to avoid the spread of weeds and pests in event of unforeseen circumstances. If spatial connectivity is not possible, proximity between patches of vegetation should be increased through the incorporation of stepping stones;
- (2) A heterogeneous network of different land cover type classes, including patches of indigenous and exotic vegetation for providing diverse habitats and food sources year-round for different species in the face of different circumstances;
- (3) Topographically diverse patches of vegetation to provide diverse micro-climatic conditions in the face of impacts imposed by rapid urbanisation and climate change;
- (4) A combination of compact and elongated patches of vegetation to meet the needs of wildlife species and people. Compact patches provide wildlife species with suitable habitats. While elongated patches play an important role in mitigating the ill effects of urban heat islands, both types of patches can supplement each other. For example, elongated patches of vegetation can be used by wildlife species as stepping-stones or additional sources of food. Likewise, compact patches can simultaneously contribute to a higher rate of evapotranspiration in order to help people and cities adapt to urban heat island effects (cf. Table 2).

Although globally accepted, the above-mentioned spatially-explicit patterns should not be considered as a panacea to address biodiversity issues in cities. In addition, they should not be used haphazardly. Instead, planning for biodiversity in cities must take *local realities on the ground* into particular consideration when using these patterns in order to maximise the odds of success. A considerate site- and species-specific attitude is therefore needed to use the spatially-explicit patterns.

Table 2: The role of landscape pattern composition and configuration in support of biodiversity in urban environments and spatially-explicit recommendations for landscape architecture

Components	Urban biodiversity targets (spatial requirements of wildlife species)	Spatially-explicit recommendations for landscape architecture
Indigenouness	<ul style="list-style-type: none">• Indigenous plant species as an essential for ecological integrity;• Patches of exotic vegetation as supplementary habitats, foraging sources, stepping stones, and buffer zones.	<ul style="list-style-type: none">• Increase the percentage of indigenous land cover type classes on an urban scale through re-vegetation, landscape restoration practices, and natural regeneration processes specifically in abandoned sites, ecologically damaged patches of vegetation, and brownfields;• Encourage people to plant indigenous species in their properties including private gardens and house backyards;• Utilise exotic species for the benefit of fauna, if they are not invasive or ecologically detrimental.
Land cover heterogeneity	<ul style="list-style-type: none">• Diversity of land cover type classes as an index of habitat diversity and thereby urban biodiversity over time.	<ul style="list-style-type: none">• Increase the number of land cover type classes on an urban scale though re-vegetation, landscape restoration practices, and natural regeneration processes;• Increase plant diversity on a micro-scale, wherever possible including along walkways, around golf courses, in playgrounds, and within cemeteries.
Land surface perviousness	<ul style="list-style-type: none">• High land surface perviousness as an index of urban wildlife diversity and abundance, more specifically birds, reptiles, and invertebrates.	<ul style="list-style-type: none">• Minimise paved areas in urban parklands, gardens, zoos, and other public green spaces;• De-pave coastal zones and increase green space coverage in urban parking lots;• Follow a more compact urban development to minimise imperviousness on a landscape level.
Patch size	<ul style="list-style-type: none">• Larger patches of vegetation as a pool for a greater number of wildlife species over time.	<ul style="list-style-type: none">• Protect large patches of vegetation from urban development and at the same time, preserve small patches of vegetation where required to provide stepping-stones or supplementary food sources between large patches to facilitate species movement.
Connectivity and proximity	<ul style="list-style-type: none">• Connected patches of vegetation as corridors for species movement;• Isolated patches of vegetation as a barrier in the face of pest and weed dispersal (i.e. predator fauna and exotic flora), as well as climatic events).	<ul style="list-style-type: none">• Plan and design green and blue corridors to connect patches of vegetation together and facilitate species movement between habitats and at the same time, allocate a number of isolated patches of vegetation to provide weed- and pest-free urban wildlife habitats in event of unforeseen circumstances.
Edge density/contrast	<ul style="list-style-type: none">• Low edge density as a strategy for reducing edge effects;• High edge contrast as a barrier in front of disease spread, pest and weed dispersal, and wildfire;• Low edge contrast as a strategy for facilitating species movement.	<ul style="list-style-type: none">• Use more compact forms when designing patches of vegetation to minimise edge density;• Minimise edge contrast through allocating similar land cover and land use activities where species movement between patches of vegetation is ecologically critical;• Maximise edge contrast through allocating dissimilar land cover and land use activities where there is a risk of disease, pest, and weed spread from one patch to another.
Landform diversity	<ul style="list-style-type: none">• Topographical diversity as a great potential for species colonisation and concentration in the event of unforeseen climate change impacts such as intensified winds, sea level rise, and cold/heat waves.	<ul style="list-style-type: none">• Avoid the modification of landform and reserve strategically important aspects of slopes as wildlife habitats and protect them from future urban development.
Shape complexity	<ul style="list-style-type: none">• Compact patches of vegetation as a medium of reducing edge effects and at the same time, increasing patch core area.	<ul style="list-style-type: none">• Use more compact forms to design new patches of vegetation.• Reconfigure the shape of patches of vegetation through plantation projects, landscape restoration practices, and natural regeneration processes, where possible.

5. Conclusion

Biodiversity loss in urban environments has already been initiated as a result of anthropogenic development; climate change is also estimated to accelerate this process. There is worldwide evidence to verify that biodiversity in urban environments, like elsewhere, depends profoundly on landscape pattern composition and configuration. If managed appropriately, the eight most important components of landscape pattern constructing spatial characteristics of urban wildlife habitats can play a key role in the presence, richness, and abundance of biodiversity in the long run despite the presence of anthropogenic development and climate change.

In-depth understanding of the role of the most important components of landscape pattern in support of wildlife species will inform the landscape architecture discipline in a way that supports biodiversity in cities, where conventionally there is not enough land, funding, and/or water to be dedicated to urban green spaces. In such situations, it is the composition and configuration of landscape patterns that contribute to urban biodiversity, not necessarily conventional development of green spaces or increasing the percentage of green space per capita.

Spatial ecology of wildlife species in urban environments is therefore essential knowledge to be developed and incorporated into the landscape architecture discipline, more than ever before, in order for urban policy makers to provide a robust basis for making informed decisions on land-use allocation and land cover conversion. This will be an initial step to help cities move from the notion of *urban development vs. habitat preservation* to *urban development with habitat conservation*.

Further research is required to address the current gaps in this field of landscape architecture research. The main research priorities include, but are not limited to, the following topics:

- (1) Given that the components of landscape pattern are not equally important to wildlife species, both global and local survey studies are required to rank the most important components of landscape pattern based upon empirical experiences gained through long-term research and practice by subject-matter experts from different parts of the world in order to provide a picture of individual importance of each component of landscape pattern in relation to others. In addition to this, such studies need to address areas of conflict between what wildlife species and humans require in terms of spatial patterning of patches of vegetation in urban environments.
- (2) Cities that have been established at ecosystem junctions (Alvey, 2006) should be identified and selected as research sites to undertake spatial analysis of landscape pattern composition and configuration in relation to the existing ecological processes in order to examine if current landscape patterns have the potentials to safeguard urban wildlife species against the local impacts of climate change and rapid urbanisation. Such research requires a reliable dataset of local information including climatic and environmental data as well as empirical information about the spatial ecology of keystone species present in the selected cities in order for landscape analysts to interpret landscape patterns in relation to the species' behaviours in space and time. The outputs are likely to reveal opportunities for the landscape architecture discipline to answer the question of what an optimised landscape pattern is to support wildlife species in the face of climate change impacts in urban environments.

References

- Ackerly, D. D., Loarie, S. R., Cornwell, W. K., Weiss, S. B., Hamilton, H., Branciforte, R., & Kraft, N. J. B. (2010). The geography of climate change: implications for conservation biogeography. *Diversity and Distributions*, 16(3), 476-487.
- Ahern, J. (1995). Greenways as a planning strategy. *Landscape and urban planning*, 33(1), 131-155.
- Aida, N., Sasidhran, S., Kamarudin, N., Aziz, N., Puan, C. L., & Azhar, B. (2016). Woody trees, green space and park size improve avian biodiversity in urban landscapes of Peninsular Malaysia. *Ecological Indicators*, 69, 176-183.
- Alberti, M. (2005). The effects of urban patterns on ecosystem function. *International regional science review*, 28(2), 168-192.
- Alvey, A. A. (2006). Promoting and preserving biodiversity in the urban forest. *Urban Forestry & Urban Greening*, 5(4), 195-201.

- Andren, H. (1994). Effects of habitat fragmentation on birds and mammals in landscapes with different proportions of suitable habitat: a review. *Oikos*, 355-366.
- Aronson, M. F., La Sorte, F. A., Nilon, C. H., Katti, M., Goddard, M. A., Lepczyk, C. A., ... & Dobbs, C. (2014). A global analysis of the impacts of urbanization on bird and plant diversity reveals key anthropogenic drivers. *Proceedings of the Royal Society of London B: Biological Sciences*, 281(1780), 20133330.
- Asgarian, A., Amiri, B. J., & Sakieh, Y. (2015). Assessing the effect of green cover spatial patterns on urban land surface temperature using landscape metrics approach. *Urban Ecosystems*, 18(1), 209-222.
- Ausseil, A. G., Dymond, J. R., Kirschbaum, M. U. F., Andrew, R. M., & Parfitt, R. L. (2013). Assessment of multiple ecosystem services in New Zealand at the catchment scale. *Environmental modelling & software*, 43, 37-48.
- Balvanera, P., Pfisterer, A. B., Buchmann, N., He, J. S., Nakashizuka, T., Raffaelli, D., & Schmid, B. (2006). Quantifying the evidence for biodiversity effects on ecosystem functioning and services. *Ecology letters*, 9(10), 1146-1156.
- Barnes, T. G. (2000). *Landscape ecology and ecosystems management*. University of Kentucky College of Agriculture, Cooperative Extension Service. Frankfort: Kentucky State University.
- Bashford, A. (2014). *Global population: history, geopolitics, and life on earth*. Columbia University Press, New York.
- Bell, S. (2001). Landscape pattern, perception and visualisation in the visual management of forests. *Landscape and Urban Planning*, 54(1), 201-211.
- Bennett, A. F. (1999). Linkages in the landscape: the role of corridors and connectivity in wildlife conservation (No. 1). *Lucn*.
- Botequilha Leitão, A. & Ahern, J. (2002). Applying landscape ecological concepts and metrics in sustainable landscape planning. *Landscape and Urban Planning*, 59(2), pp. 65-93.
- Botequilha Leitão, A., Miller, J., Ahern, J. & McGarigal, K. (2006). *Measuring landscapes: A planner's handbook*. Island Press, Washington.
- Bräuniger, C., Knapp, S., Kühn, I., & Klotz, S. (2010). Testing taxonomic and landscape surrogates for biodiversity in an urban setting. *Landscape and Urban Planning*, 97(4), 283-295.
- Bridgewater, P. B. (1988). Biodiversity and landscape. *Earth-Science Reviews*, 25(5), 485-491.
- Calder, J. L., Cumming, G. S., Maciejewski, K., & Oschadleus, H. D. (2015). Urban land use does not limit weaver bird movements between wetlands in Cape Town, South Africa. *Biological Conservation*, 187, 230-239.
- Carbó-Ramírez, P., & Zuria, I. (2011). The value of small urban greenspaces for birds in a Mexican city. *Landscape and Urban Planning*, 100(3), 213-222.
- Cardinale, B. J., Srivastava, D. S., Duffy, J. E., Wright, J. P., Downing, A. L., Sankaran, M., & Jouseau, C. (2006). Effects of biodiversity on the functioning of trophic groups and ecosystems. *Nature*, 443(7114), 989-992.
- Chong, K. Y., Teo, S., Kurukulasuriya, B., Chung, Y. F., Rajathurai, S., & Tan, H. T. W. (2014). Not all green is as good: Different effects of the natural and cultivated components of urban vegetation on bird and butterfly diversity. *Biological Conservation*, 171, 299-309.
- Collinge, S. K. (1996). Ecological consequences of habitat fragmentation: implications for landscape architecture and planning. *Landscape and urban planning*, 36(1), 59-77.
- Cornelis, J., & Hermy, M. (2004). Biodiversity relationships in urban and suburban parks in Flanders. *Landscape and Urban Planning*, 69(4), 385-401.
- Davies-Colley, R. J., Payne, G. W., & Van Elswijk, M. (2000). Microclimate gradients across a forest edge. *New Zealand Journal of Ecology*, 24, 111-121.
- de Toledo, M. C. B., Donatelli, R. J., & Batista, G. T. (2012). Relation between green spaces and bird community structure in an urban area in Southeast Brazil. *Urban Ecosystems*, 15(1), 111-131.
- Dramstad, W. E., Olson, J. D. & Forman, R. T. T. (1996) *Landscape ecology principles in landscape architecture and land-use planning*. Harvard University Graduate School of Design and Island Press, Washington.
- Duelli, P. (1997). Biodiversity evaluation in agricultural landscapes: an approach at two different scales. *Agriculture, ecosystems & environment*, 62(2), 81-91.
- Duffy, J. E. (2008). Why biodiversity is important to the functioning of real-world ecosystems. *Frontiers in Ecology and the Environment*, 7(8), 437-444.
- Duffy, J. E., Cardinale, B. J., France, K. E., McIntyre, P. B., Thébault, E., & Loreau, M. (2007). The functional role of biodiversity in ecosystems: incorporating trophic complexity. *Ecology letters*, 10(6), 522-538.
- ELI - Environmental Law Institute (2003) *Conservation thresholds for land use planners*. Environmental Law Institute, Washington DC.
- Evans, K. L., Newson, S. E., & Gaston, K. J. (2009). Habitat influences on urban avian assemblages. *Ibis*, 151(1), 19-39.
- Fabos, J. G. (2004). Greenway planning in the United States: its origins and recent case studies. *Landscape and urban planning*, 68(2), 321-342.
- Farina, A. (2006). *Principles and methods in landscape ecology. Toward a science of landscape*. Landscape Series, Vol. 3. Springer, Dordrecht.

- Ferenc, M., Sedláček, O., & Fuchs, R. (2014). How to improve urban greenspace for woodland birds: site and local-scale determinants of bird species richness. *Urban Ecosystems*, 17(2), 625-640.
- Fernández-Jurícic, E. & Jokimaki, J. (2001). A habitat island approach to conserving birds in urban landscapes: case studies from southern and northern Europe. *Biodiversity and Conservation* 10: 2023-2043.
- Fernández-Jurícic, E. (2001). Avian spatial segregation at edges and interiors of urban parks in Madrid, Spain. *Biodiversity Conservation*. 10, 1303-1316.
- Feyisa, G. L., Dons, K., & Meilby, H. (2014). Efficiency of parks in mitigating urban heat island effect: An example from Addis Ababa. *Landscape and Urban Planning*, 123, 87-95.
- Fischer, J., Lindenmayer, D.B., Hobbs, R. (2009). Landscape pattern and biodiversity. pp. 431–437. In Levin S. A. (ed.), *The Princeton guide to ecology*, New Jersey.
- Fontana, S., Sattler, T., Bontadina, F., & Moretti, M. (2011). How to manage the urban green to improve bird diversity and community structure. *Landscape and Urban Planning*, 101(3), 278-285.
- Forman, R. T. T. (1995). *Land mosaics: The ecology of landscapes and regions*. Cambridge University Press, Cambridge.
- Forman, R. T. T., & Godron, M. (1986). *Landscape ecology*. 619 pp. Jhon Wiley & Sons, New York.
- Forman, R. T., & Collinge, S. K. (1997). Nature conserved in changing landscapes with and without spatial planning. *Landscape and urban planning*, 37(1), 129-135.
- Forman, R. T., & Godron, M. (1981). Patches and structural components for a landscape ecology. *BioScience*, 31(10), 733-740.
- Forman, R. T., Galli, A. E., & Leck, C. F. (1976). Forest size and avian diversity in New Jersey woodlots with some land use implications. *Oecologia*, 26(1), 1-8.
- Garden, J. G., McAlpine, C. A., & Possingham, H. P. (2010). Multi-scaled habitat considerations for conserving urban biodiversity: native reptiles and small mammals in Brisbane, Australia. *Landscape Ecology*, 25(7), 1013-1028.
- Grimm, N. B., Faeth, S. H., Golubiewski, N. E., Redman, C. L., Wu, J., Bai, X., & Briggs, J. M. (2008). Global change and the ecology of cities. *Science*, 319(5864), 756-760.
- Haddad, N. (2009). Principles of reserve design. pp. 529-537. In Levin S. A. (ed.), *The Princeton guide to ecology*, New Jersey.
- Hahs, A. K., McDonnell, M. J., McCarthy, M. A., Vesk, P. A., Corlett, R. T., Norton, B. A., ... & Williams, N. S. (2009). A global synthesis of plant extinction rates in urban areas. *Ecology Letters*, 12(11), 1165-1173.
- Halpin, P. N. (1997). Global climate change and natural-area protection: management responses and research directions. *Ecological Applications*, 7(3), 828-843.
- Hector, A., & Bagchi, R. (2007). Biodiversity and ecosystem multifunctionality. *Nature*, 448(7150), 188-190.
- Heller, N. E., & Zavaleta, E. S. (2009). Biodiversity management in the face of climate change: a review of 22 years of recommendations. *Biological conservation*, 142(1), 14-32.
- Honnay, O., Piessens, K., Van Landuyt, W., Hermy, M., & Gulinck, H. (2003). Satellite based land use and landscape complexity indices as predictors for regional plant species diversity. *Landscape and urban planning*, 63(4), 241-250.
- Hooper, D. U., Adair, E. C., Cardinale, B. J., Byrnes, J. E., Hungate, B. A., Matulich, K. L., ... & O'Connor, M. I. (2012). A global synthesis reveals biodiversity loss as a major driver of ecosystem change. *Nature*, 486(7401), 105-108.
- Ikin, K., Beaty, R. M., Lindenmayer, D. B., Knight, E., Fischer, J., & Manning, A. D. (2013). Pocket parks in a compact city: how do birds respond to increasing residential density?. *Landscape ecology*, 28(1), 45-56.
- Imai, H., & Nakashizuka, T. (2010). Environmental factors affecting the composition and diversity of avian community in mid-to late breeding season in urban parks and green spaces. *Landscape and Urban Planning*, 96(3), 183-194.
- IPCC, (1995). *Scientific Assessments of Climate Change. The Policymaker's Summary of Working Group 1 to the Intergovernmental Panel on Climate Change*, WMO/UNEP
- IPCC, (2014). *Summary for Policymakers. In: Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* [Edenhofer, O., R. Pichs-Madruga, Y. Sokona, E. Farahani, S. Kadner, K. Seyboth, A. Adler, I. Baum, S. Brunner, P. Eickemeier, B. Kriemann, J. Savolainen, S. Schlömer, C. von Stechow, T. Zwickel and J.C. Minx (eds.)]. Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA.
- Isbell, F., Calcagno, V., Hector, A., Connolly, J., Harpole, W. S., Reich, P. B., ... & Weigelt, A. (2011). High plant diversity is needed to maintain ecosystem services. *Nature*, 477(7363), 199-202.
- Jasmani, Z., Ravn, H. P., & van den Bosch, C. C. K. (2017). The influence of small urban parks characteristics on bird diversity: A case study of Petaling Jaya, Malaysia. *Urban Ecosystems*, 20(1), 227-243.
- Kang, W., Minor, E. S., Park, C. R., & Lee, D. (2015). Effects of habitat structure, human disturbance, and habitat connectivity on urban forest bird communities. *Urban Ecosystems*, 18(3), 857-870.
- Khera, N., Mehta, V., & Sabata, B. C. (2009). Interrelationship of birds and habitat features in urban greenspaces in Delhi, India. *Urban Forestry & Urban Greening*, 8(3), 187-196.
- Kong, F., Yin, H., Wang, C., Cavan, G., & James, P. (2014). A satellite image-based analysis of factors contributing to the green-space cool island intensity on a city scale. *Urban Forestry & Urban Greening*, 13(4), 846-853.

- Lefcheck, J. S., & Duffy, J. E. (2015). Multitrophic functional diversity predicts ecosystem functioning in experimental assemblages of estuarine consumers. *Ecology*, 96(11), 2973-2983.
- Leveau, L. M., & Leveau, C. M. (2016). Does urbanization affect the seasonal dynamics of bird communities in urban parks?. *Urban ecosystems*, 19(2), 631-647.
- Lizée, M. H., Manel, S., Mauffrey, J. F., Tatoni, T., & Deschamps-Cottin, M. (2012). Matrix configuration and patch isolation influences override the species–area relationship for urban butterfly communities. *Landscape ecology*, 27(2), 159-169.
- MacArthur, R., & Wilson, E. (1967). *The theory of biogeography*. Princeton University Press, New Jersey.
- MacLeod, C. J., Tinkler, G., Gormley, A. M., & Spurr, E. B. (2012). Measuring occupancy for an iconic bird species in urban parks. *New Zealand Journal of Ecology*, 36(3), 1-10.
- Maimaitiyiming, M., Ghulam, A., Tiyp, T., Pla, F., Latorre-Carmona, P., Halik, Ü., ... & Caetano, M. (2014). Effects of green space spatial pattern on land surface temperature: Implications for sustainable urban planning and climate change adaptation. *ISPRS Journal of Photogrammetry and Remote Sensing*, 89, 59-66.
- Markham, A., Dudley, N., & Stolton, S. (1993). Some like it hot: climate change, biodiversity and the survival of species.
- Mason, J., Moorman, C., Hess, G., & Sinclair, K. (2007). Designing suburban greenways to provide habitat for forest-breeding birds. *Landscape and Urban Planning*, 80(1), 153-164.
- McClure, C. J., Korte, A. C., Heath, J. A., & Barber, J. R. (2015). Pavement and riparian forest shape the bird community along an urban river corridor. *Global Ecology and Conservation*, 4, 291-310.
- McGarigal, K., & Cushman, S.A. (2002). Comparative evaluation of experimental approaches to the study of habitat fragmentation effects. *Ecological applications*, 12(2), 335-345.
- McHale, M. R., Hall, S. J., Majumdar, A., & Grimm, N. B. (2017). Carbon lost and carbon gained: a study of vegetation and carbon trade-offs among diverse land uses in Phoenix, Arizona. *Ecological Applications*, 27(2), 644-661.
- McKinney, M. L. (2002). Urbanization, Biodiversity, and Conservation. *BioScience*, 52(10), 883-890.
- Meurk, C. D., & Hall, G. M. (2006). Options for enhancing forest biodiversity across New Zealand's managed landscapes based on ecosystem modelling and spatial design. *New Zealand Journal of Ecology*, (30)1: 131-146.
- Meurk, C. D., & Swaffield, S. R. (2000). A landscape ecological framework for indigenous regeneration in rural New Zealand-Aotearoa. *Landscape and Urban Planning*, 50(1), 129-144.
- Millennium Ecosystem Assessment, (2005). *Current state and trends*. World Resources Institute, Washington DC.
- Ministry for Primary Industries, (2015). *Look-up Tables for Forestry in the Emissions Trading Scheme*. Wellington, New Zealand.
- Moorcroft, P. R. (2009). Biodiversity patterns in managed and natural landscapes. pp. 445-457. In Levin S. A. (ed.), *The Princeton guide to ecology*, New Jersey.
- Morrison, M. L., Marcot, B., & Mannan, W. (2012). *Wildlife-habitat relationships: concepts and applications*. Island Press, Washington.
- Müller, N., & Kamada, M. (2011). URBIO: an introduction to the International Network in Urban Biodiversity and Design. *Landscape and ecological engineering*, 7(1), 1-8.
- Murcia, C. (1995). Edge effects in fragmented forests: implications for conservation. *Trends in ecology & evolution*, 10(2), 58-62.
- Murgui, E. (2009). Influence of urban landscape structure on bird fauna: a case study across seasons in the city of Valencia (Spain). *Urban ecosystems*, 12(3), 249.
- Naveh, Z. (1994). From Biodiversity to Ecodiversity: A Landscape Ecology Approach to Conservation and Restoration. *Restoration Ecology*, 2(3), 180-189.
- Nielsen, A. B., van den Bosch, M., Maruthaveeran, S., & van den Bosch, C. K. (2014). Species richness in urban parks and its drivers: a review of empirical evidence. *Urban ecosystems*, 17(1), 305-327.
- Noss, R. F. (2001). Beyond Kyoto: forest management in a time of rapid climate change. *Conservation biology*, 15(3), 578-590.
- Noss, R. F., & Cooperrider, A. (1994). *Saving nature's legacy: protecting and restoring biodiversity*. Island Press, Washington.
- Oliver, A. J., Hong-Wa, C., Devonshire, J., Olea, K. R., Rivas, G. F., & Gahl, M. K. (2011). Avifauna richness enhanced in large, isolated urban parks. *Landscape and Urban Planning*, 102(4), 215-225.
- O'Neill, R. V., Milne, B. T., Turner, M. G., & Gardner, R. H. (1988). Resource utilization scales and landscape pattern. *Landscape Ecology*, 2(1), 63-69.
- Opdam, P., Steingröver, E., van Rooij, S. (2006). Ecological networks: A spatial concept for multi-actor planning of sustainable landscapes. *Landscape and Urban Planning*, 75: 322-332.
- Pacheco, R., & Vasconcelos, H. L. (2007). Invertebrate conservation in urban areas: ants in the Brazilian Cerrado. *Landscape and Urban Planning*, 81(3), 193-199.
- Paker, Y., Yom-Tov, Y., Alon-Mozes, T., & Barnea, A. (2014). The effect of plant richness and urban garden structure on bird species richness, diversity and community structure. *Landscape and urban planning*, 122, 186-195.

- Palmisano, G. O., Govindan, K., Loisi, R. V., Dal Sasso, P., & Roma, R. (2016). Greenways for rural sustainable development: An integration between geographic information systems and group analytic hierarchy process. *Land Use Policy*, 50, 429-440.
- Park, J. H., & Cho, G. H. (2016). Examining the Association between Physical Characteristics of Green Space and Land Surface Temperature: A Case Study of Ulsan, Korea. *Sustainability*, 8(8), 777.
- Parnesan, C. (2006). Ecological and evolutionary responses to recent climate change. *Annual Review of Ecology, Evolution, and Systematics*, 37, 637-669.
- Pasari, J. R., Levi, T., Zavaleta, E. S., & Tilman, D. (2013). Several scales of biodiversity affect ecosystem multifunctionality. *Proceedings of the National Academy of Sciences*, 110(25), 10219-10222.
- Pedersen Zari, M. (2015). Ecosystem services analysis: Mimicking ecosystem services for regenerative urban design. *International Journal of Sustainable Built Environment*, 4, 145-157.
- Peel, M. C., Finlayson, B. L., & McMahon, T. A. (2007). Updated world map of the Köppen-Geiger climate classification. *Hydrology and Earth System Sciences Discussions*, 4(2), 439-473.
- Pernetta, J. L., Elder, R., & D Humphrey, S. (1994). Impacts of climate change on ecosystems and species: Implications for protected areas. In 4. World Congress on National Parks and Protected Areas Ene 1992 Caracas (Venezuela) (No. 333.95 I34). IUCN, Gland (Suiza). Marine and Coastal Areas Programme.
- Platt, A., & Lill, A. (2006). Composition and conservation value of bird assemblages of urban 'habitat islands': do pedestrian traffic and landscape variables exert an influence?. *Urban Ecosystems*, 9(2), 83-97.
- Port Stephens Council. (2002). Port Stephens Council Comprehensive Koala Plan of Management (CKPoM) – June 2002. Prepared by Port Stephens Council with the Australian Koala Foundation.
- Pulliam, H. R. (1988). Sources, sinks, and population regulation. *American naturalist*, 132, 652-661.
- Rastandeh, A., Pedersen Zari, M., K. Brown, D., & Vale, R. (2017). Utilising exotic flora in support of urban indigenous biodiversity: lessons for landscape architecture. *Landscape Research*, 1-13.
- Rega, C. C., Nilon, C. H., & Warren, P. S. (2015). Avian abundance patterns in relation to the distribution of small urban greenspaces. *Journal of Urban Planning and development*, 141(3), A4015002.
- Reside, A. E., Welbergen, J. A., Phillips, B. L., Wardell-Johnson, G. W., Keppel, G., Ferrier, S., ... & VanDerWal, J. (2014). Characteristics of climate change refugia for Australian biodiversity. *Austral Ecology*, 39(8), 887-897.
- Rorison, I. H., Sutton, F., & Hunt, R. (1986). Local climate, topography and plant growth in Lathkill Dale NNR. I. A twelve-year summary of solar radiation and temperature. *Plant, Cell & Environment*, 9(1), 49-56.
- Sandström, U. G., Angelstam, P., & Mikusiński, G. (2006). Ecological diversity of birds in relation to the structure of urban green space. *Landscape and urban planning*, 77(1), 39-53.
- Sanz, V., & Caula, S. (2015). Assessing bird assemblages along an urban gradient in a Caribbean island (Margarita, Venezuela). *Urban ecosystems*, 18(3), 729-746.
- Savard, J. P. L., Clergeau, P., & Mennechez, G. (2000). Biodiversity concepts and urban ecosystems. *Landscape and urban planning*, 48(3), 131-142.
- Setälä, H. M., Francini, G., Allen, J. A., Hui, N., Jumpponen, A., & Kotze, D. J. (2016). Vegetation Type and Age Drive Changes in Soil Properties, Nitrogen, and Carbon Sequestration in Urban Parks under Cold Climate. *Frontiers in Ecology and Evolution*, 4, 93.
- Shaffer, M. L. (1981). Minimum population sizes for species conservation. *Biological Science*, 31(2), 131-134.
- Shwartz, A., Shirley, S., & Kark, S. (2008). How do habitat variability and management regime shape the spatial heterogeneity of birds within a large Mediterranean urban park?. *Landscape and Urban Planning*, 84(3), 219-229.
- Sing, K. W., Jusoh, W. F., Hashim, N. R., & Wilson, J. J. (2016). Urban parks: refuges for tropical butterflies in Southeast Asia?. *Urban ecosystems*, 19(3), 1131-1147.
- Smith, P. G. (2007). Characteristics of urban natural areas influencing winter bird use in southern Ontario, Canada. *Environmental management*, 39(3), 338-352.
- Soule, M.E. (1991). Land use planning and wildlife maintenance: guidelines for conserving wildlife in an urban landscape. *Journal of the American Planning Association*, 57(3): 313-323.
- Strohbach, M. W., Lerman, S. B., & Warren, P. S. (2013). Are small greening areas enhancing bird diversity? Insights from community-driven greening projects in Boston. *Landscape and Urban Planning*, 114, 69-79.
- Sullivan, J. J., Meurk, C., Whaley, K. J., & Simcock, R. (2009). Restoring native ecosystems in urban Auckland: urban soils, isolation, and weeds as impediments to forest establishment. *New Zealand Journal of Ecology*, 33(1): 60-71.
- Suri, J., Anderson, P. M., Charles-Dominique, T., Hellard, E., & Cumming, G. S. (2017). More than just a corridor: A suburban river catchment enhances bird functional diversity. *Landscape and Urban Planning*, 157, 331-342.
- Teng, M., Wu, C., Zhou, Z., Lord, E., & Zheng, Z. (2011). Multipurpose greenway planning for changing cities: A framework integrating priorities and a least-cost path model. *Landscape and urban planning*, 103(1), 1-14.
- Threlfall, C. G., Williams, N. S., Hahs, A. K., & Livesley, S. J. (2016). Approaches to urban vegetation management and the impacts on urban bird and bat assemblages. *Landscape and Urban Planning*, 153, 28-39.

- Tilman, D., Isbell, F., & Cowles, J. M. (2014). Biodiversity and ecosystem functioning. *Annual Review of Ecology, Evolution, and Systematics*, 45(1), 471.
- Toccolini, A., Fumagalli, N., & Senes, G. (2006). Greenways planning in Italy: the Lambro River Valley greenways system. *Landscape and urban planning*, 76(1), 98-111.
- Turner, M. G. (1989) Landscape ecology: The effect of pattern on process. *Annual Review of Ecology and Systematics*, 20, pp. 97-171.
- Turner, T. (2006). Greenway planning in Britain: recent work and future plans. *Landscape and urban planning*, 76(1), 240-251.
- Tzoulas, K., Korpela, K., Venn, S., Yli-Pelkonen, V., Kaźmierczak, A., Niemela, J., & James, P. (2007). Promoting ecosystem and human health in urban areas using Green Infrastructure: A literature review. *Landscape and urban planning*, 81(3), 167-178.
- United Nations - Department of Economic and Social Affairs, Population Division (2014). World urbanization prospects: the 2014 revision, Highlights (ST/ESA/SER.A/352).
- United Nations - Department of Economic and Social Affairs, Population Division (2008). World urbanization prospects: the 2007 revision. CD-ROM Edition, data in digital form (POP/DB/WUP/Rev.2007).
- Vallejo, B. M., Aloy, A. B., & Ong, P. S. (2009). The distribution, abundance and diversity of birds in Manila's last greenspaces. *Landscape and Urban Planning*, 89(3), 75-85.
- van Heezik, Y., Freeman, C., Porter, S., & Dickinson, K. J. (2013). Garden size, householder knowledge, and socio-economic status influence plant and bird diversity at the scale of individual gardens. *Ecosystems*, 16(8), 1442-1454.
- van Heezik, Y., Smyth, A., & Mathieu, R. (2008). Diversity of native and exotic birds across an urban gradient in a New Zealand city. *Landscape and urban planning*, 87(3), 223-232.
- Villegas, M., & Garitano-Zavala, Á. (2010). Bird community responses to different urban conditions in La Paz, Bolivia. *Urban Ecosystems*, 13(3), 375-391.
- Walmsley, A. (2006). Greenways: multiplying and diversifying in the 21st century. *Landscape and urban planning*, 76(1), 252-290.
- Walz, U., & Syrbe, R. U. (2013). Linking landscape structure and biodiversity. *Ecological indicators*, 31, 1-5.
- Young, A., & Mitchell, N. (1994). Microclimate and vegetation edge effects in a fragmented podocarp-broadleaf forest in New Zealand. *Biological Conservation*, 67(1), 63-72.
- Yuan, B. A. O. D. O. N. G., & Lu, C. (2016). Effects of urbanization on bird diversity: a case study in Yizhou, Guangxi Province, China. *Asia Life Sciences*, 25, 79-96.
- Zhou, D., & Chu, L. M. (2012). How would size, age, human disturbance, and vegetation structure affect bird communities of urban parks in different seasons?. *Journal of Ornithology*, 153(4), 1101-1112.
- Zhou, W., Huang, G., & Cadenasso, M. L. (2011). Does spatial configuration matter? Understanding the effects of land cover pattern on land surface temperature in urban landscapes. *Landscape and Urban Planning*, 102(1), 54-63.